

University of Groningen

**Glucocorticoid enhancement of memory requires arousal-induced noradrenergic activation in the basolateral amygdala**

Roozendaal, B; Okuda, S; van der Zee, E A; McGaugh, James L.

*Published in:*

Proceedings of the National Academy of Sciences of the United States of America

*DOI:*

[10.1073/pnas.0601874103](https://doi.org/10.1073/pnas.0601874103)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2006

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Roozendaal, B., Okuda, S., van der Zee, E. A., & McGaugh, J. L. (2006). Glucocorticoid enhancement of memory requires arousal-induced noradrenergic activation in the basolateral amygdala. *Proceedings of the National Academy of Sciences of the United States of America*, 103(17), 6741-6746.  
<https://doi.org/10.1073/pnas.0601874103>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# Glucocorticoid enhancement of memory requires arousal-induced noradrenergic activation in the basolateral amygdala

Roozendaal *et al.* 10.1073/pnas.0601874103.

## Supporting Tables

Table 2. Total object exploration time

Habitation	Drug treatment	Training	Retention
No Prior Habituation	Vehicle	23.4 ± 2.0 (11)	17.6 ± 1.6
	Cort (0.3 mg/kg)	20.7 ± 2.1 (12)	19.1 ± 1.4
	Cort (1.0 mg/kg)	27.5 ± 2.6 (10)	19.9 ± 1.6
	Cort (3.0 mg/kg)	26.9 ± 3.0 (10)	19.1 ± 2.3
	Propranolol (3.0 mg/kg)	27.7 ± 2.0 (11)	18.3 ± 1.7
	Cort (0.3 mg/kg) + propranolol	27.1 ± 2.9 (8)	14.6 ± 2.0
	Cort (1.0 mg/kg) + propranolol	23.2 ± 1.3 (12)	15.5 ± 0.7
	Cort (3.0 mg/kg) + propranolol	22.7 ± 1.4 (11)	19.2 ± 1.8
	One-way ANOVA	$F_{7,77}=1.62; P=0.14$	$F_{7,77}=1.26; P=0.28$
Prior Habituation	Vehicle	30.5 ± 2.1 (16)	25.1 ± 3.2
	Cort (0.3 mg/kg)	33.8 ± 4.0 (12)	24.2 ± 2.5
	Cort (1.0 mg/kg)	31.7 ± 2.3 (17)	21.7 ± 1.6
	Cort (3.0 mg/kg)	33.7 ± 3.7 (12)	21.2 ± 1.4

	Yohimbine (0.3 mg/kg)	32.2 ± 2,6 (10)	19.8 ± 1.9
	Cort (0.3 mg/kg) + yohimbine	30.2 ± 1.8 (9)	20.5 ± 2.6
	Cort (1.0 mg/kg) + yohimbine	27.4 ± 2.6 (12)	16.6 ± 1.5
	Cort (3.0 mg/kg) + yohimbine	32.6 ± 2.0 (9)	18.5 ± 1.3
	One-way ANOVA	$F_{7,89}=0.56; P=0.79$	$F_{7,89}=1.62; P=0.14$

Total time spent exploring the two objects (two identical objects for the training trial, and a familiar and novel object for the test trial), expressed as mean ± SEM in seconds. The numbers of animals per group are shown in parentheses. Cort, corticosterone.

**Table 3. Total object exploration time**

Basolateral amygdala	Systemic	Training	Retention
Saline	Vehicle	19.7 ± 2.0 (10)	17.9 ± 1.7
Saline	Cort (0.3 mg/kg)	20.1 ± 1.5 (10)	16.3 ± 1.2
Saline	Cort (1.0 mg/kg)	24.3 ± 1.7 (10)	20.8 ± 2.4
Saline	Cort (3.0 mg/kg)	19.7 ± 0.7 (9)	18.9 ± 2.1
Propranolol (0.5 µg)	Vehicle	22.0 ± 1.2 (8)	17.6 ± 0.9
Propranolol (0.5 µg)	Cort (0.3 mg/kg)	21.3 ± 1.6 (11)	17.6 ± 1.6
Propranolol (0.5 µg)	Cort (1.0 mg/kg)	20.3 ± 1.4 (11)	16.5 ± 1.0

Propranolol (0.5 µg)	Cort (3.0 mg/kg)	21.7 ± 1.2 (10)	18.0 ± 2.2
	One-way ANOVA	$F_{7,71}=1.08;$ $P=0.39$	$F_{7,71}=0.69;$ $P=0.68$
Hippocampus	Systemic	Training	Retention
Saline	Vehicle	25.0 ± 1.0 (10)	21.0 ± 2.0
Saline	Cort (0.3 mg/kg)	27.1 ± 2.4 (13)	19.9 ± 2.3
Saline	Cort (1.0 mg/kg)	25.2 ± 2.1 (14)	19.7 ± 1.1
Saline	Cort (3.0 mg/kg)	25.0 ± 1.6 (11)	18.6 ± 2.4
Propranolol (1.25 µg)	Vehicle	26.6 ± 1.3 (10)	19.6 ± 1.9
Propranolol (1.25 µg)	Cort (0.3 mg/kg)	27.7 ± 2.5 (11)	19.5 ± 1.8
Propranolol (1.25 µg)	Cort (1.0 mg/kg)	24.7 ± 2.3 (13)	20.5 ± 1.5
Propranolol (1.25 µg)	Cort (3.0 mg/kg)	24.3 ± 1.5 (12)	18.6 ± 1.1
	One-way ANOVA	$F_{7,86}=0.41;$ $P=0.89$	$F_{7,86}=0.20;$ $P=0.98$

Total time spent exploring the two objects (two identical objects for the training trial, and a familiar and novel object for the test trial), expressed as mean ± SEM in seconds. The numbers of animals per group are shown in parentheses. Cort, corticosterone.